



TITLE:

# ON THE NORM CONVERGENCE OF THE TROTTER-KATO PRODUCT FORMULA WITH ERROR BOUND (Spectral and Scattering Theory and Related Topics)

AUTHOR(S):

Ichinose, Takashi; Tamura, Hideo

---

CITATION:

Ichinose, Takashi ...[et al]. ON THE NORM CONVERGENCE OF THE TROTTER-KATO PRODUCT FORMULA WITH ERROR BOUND (Spectral and Scattering Theory and Related Topics). 数理解析研究所講究録 2001, 1208: 128-134

ISSUE DATE:

2001-05

URL:

<http://hdl.handle.net/2433/41064>

RIGHT:

# ON THE NORM CONVERGENCE OF THE TROTTER-KATO PRODUCT FORMULA WITH ERROR BOUND

金沢大理 一瀬 孝 (Takashi Ichinose\*)

岡山大理 田村 英男 (Hideo Tamura\*\*)

\*Department of Mathematics, Faculty of Science, Kanazawa Univ.

\*\* Department of Mathematics, Faculty of Science, Okayama Univ.

**Abstract.** The norm convergence of the Trotter-Kato product formula with error bound is shown for the semigroup generated by that operator sum of two nonnegative selfadjoint operators  $A$  and  $B$  which is selfadjoint.

## 1. Introduction and Result

It is well-known ([23], [15]; [19]) that the Trotter-Kato product formula for the selfadjoint semigroup holds in strong operator topology. Namely, when  $A$  and  $B$  are nonnegative selfadjoint operators in a Hilbert space  $\mathcal{H}$  with domains  $D[A]$  and  $D[B]$ , then

$$\text{s-lim}_{n \rightarrow \infty} (e^{-tB/2n} e^{-tA/n} e^{-tB/2n})^n = \text{s-lim}_{n \rightarrow \infty} (e^{-tA/n} e^{-tB/n})^n = e^{-tC}, \quad (1.1)$$

if  $C$  is the form sum  $A+B$  which is selfadjoint, or, in particular, if the operator sum  $A+B$  is essentially selfadjoint on  $D[A] \cap D[B]$  with  $C$  its closure. The convergence is uniform on each compact  $t$ -interval in the closed half line  $[0, \infty)$ .

The aim of this note is to briefly announce our recent results on its operator-norm convergence with error bound. In [12] we have shown

**Theorem 1.1.** *If  $A$  and  $B$  are nonnegative selfadjoint operators in  $\mathcal{H}$  with domains  $D[A]$  and  $D[B]$  and if their operator sum  $C := A+B$  is selfadjoint on  $D[C] = D[A] \cap D[B]$ , then the product formula in operator norm holds with error bound:*

$$\begin{aligned} \|(e^{-tB/2n} e^{-tA/n} e^{-tB/2n})^n - e^{-tC}\| &= O(n^{-1/2}), \\ \|(e^{-tA/n} e^{-tB/n})^n - e^{-tC}\| &= O(n^{-1/2}), \quad n \rightarrow \infty. \end{aligned} \quad (1.2)$$

The convergence is uniform on each compact  $t$ -interval in the open half line  $(0, \infty)$ , and further, if  $C$  is strictly positive, uniform on the closed half line  $[T, \infty)$  for every fixed  $T > 0$ .

One of the typical examples of such a selfadjoint operator  $C = A + B$  is the Schrödinger operator

$$H = -\frac{1}{2}\Delta + P|x|^{-1} + D|x|^2 + E|x|^{2000}$$

in  $L^2(\mathbf{R}^3)$ , where  $P$ ,  $D$  and  $E$  are nonnegative constants.

*Remark 1.1* The first result of such a norm convergence of the Trotter–Kato product formula (1.1) was proved by Rogava [20] in the abstract case under an additional condition that  $B$  is  $A$ -bounded, with error bound  $O(n^{-1/2} \log n)$ . The next was by Helffer [5] for the Schrödinger operators  $H = H_0 + V \equiv -\frac{1}{2}\Delta + V(x)$  with  $C^\infty$  nonnegative potentials  $V(x)$ , roughly speaking, growing at most of order  $O(|x|^2)$  for large  $|x|$  with error bound  $O(n^{-1})$ . Each of these two results is independent of the other.

Then under some stronger or more general conditions, several further results are obtained. As for the abstract case, a better error bound  $O(n^{-1} \log n)$  than Rogava's is obtained by Ichinose–Tamura [11] (cf. [9]) when  $B$  is  $A^\alpha$ -bounded for some  $0 < \alpha < 1$ , even though the  $B = B(t)$  may be  $t$ -dependent, and by Neidhardt–Zagrebnov [16], [17] (cf. [18]) when  $B$  is  $A$ -bounded with relative bound less than 1. As for the Schrödinger operators, a different proof to Helffer's result was obtained by Dia-Schatzman [2]. Further, more general results were proved for continuous nonnegative potentials  $V(x)$ , roughly speaking, growing of order  $O(|x|^\rho)$  for large  $|x|$  with  $\rho > 0$ , together with error bounds dependent on the power  $\rho$  (for instance, of order  $O(n^{-2/\rho})$ , if  $\rho \geq 2$ ), by Ichinose–Takanobu [6] (cf. [7]), Doumeki–Ichinose–Tamura [3], Ichinose–Tamura [10], Decombes–Dia [1] and others, although the primary purpose of most of these papers was to prove rather a norm estimate between the Kac transfer operator and its corresponding Schrödinger semigroup. The Schrödinger operators treated in [6] and [3] may even involve bounded magnetic fields  $\nabla \times A(x)$ :  $H = H_0(A) + V \equiv \frac{1}{2}(-i\nabla - A(x))^2 + V(x)$ . In [7] and [8] the relativistic Schrödinger operator was also dealt with.

It should be noted (see [4], [21]) that in all these cases of the Schrödinger operators the sum  $H = H_0 + V$  (resp.  $H = H_0(A) + V$ ) is selfadjoint on the domain  $D[H] = D[H_0] \cap D[V]$  (resp.  $D[H] = D[H_0(A)] \cap D[V]$ ).

Thus the present theorem not only extends Rogava's result, but also can extend and contain all the results mentioned above, inclusive better error bounds in some cases.

*Remark 1.2.* Unless the sum  $A + B$  is selfadjoint on  $D[A] \cap D[B]$ , the norm convergence of the Trotter–Kato product formula does not always hold, even though the sum is essentially selfadjoint there and  $B$  is  $A$ -form-bounded with relative bound less than 1. A counterexample is due to Hiroshi Tamura [22].

The theorem also holds with the exponential function  $e^{-s}$  replaced by real-valued, Borel measurable functions  $f$  and  $g$  on  $[0, \infty)$  satisfying that

$$0 \leq f(s) \leq 1, \quad f(0) = 1, \quad f'(0) = -1, \quad (1.3)$$

that for every small  $\varepsilon > 0$  there exists a positive constant  $\delta = \delta(\varepsilon) < 1$  such that

$$f(s) \leq 1 - \delta(\varepsilon), \quad s \geq \varepsilon, \quad (1.4)$$

and that, for some fixed constant  $\kappa$  with  $1 < \kappa \leq 2$ ,

$$[f]_\kappa := \sup_{s>0} s^{-\kappa} |f(s) - 1 + s| < \infty, \quad (1.5)$$

and the same for  $g$ . Of course, the functions  $f(s) = e^{-s}$  and  $f(s) = (1 + k^{-1}s)^{-k}$  with  $k > 0$  are examples of functions having these properties.

**Theorem 1.2.** *If  $3/2 \leq \kappa \leq 2$ , it holds in operator norm that*

$$\begin{aligned} \| [g(tB/2n)f(tA/n)g(tB/2n)]^n - e^{-tC} \| &= O(n^{-1/2}), \\ \| [f(tA/n)g(tB/n)]^n - e^{-tC} \| &= O(n^{-1/2}), \quad n \rightarrow \infty. \end{aligned} \quad (1.6)$$

## 2. Outline of Proof

To proving the theorem, it is crucial to show the following operator-norm version of Chernoff's theorem with error bounds. The case without error bounds was noted by Neidhardt-Zagrebnov [18].

**Lemma.** *Let  $C$  be a nonnegative selfadjoint operator in a Hilbert space  $\mathcal{H}$  and let  $\{F(t)\}_{t \geq 0}$  be a family of selfadjoint operators with  $0 \leq F(t) \leq 1$ . Define  $S_t = t^{-1}(1 - F(t))$ . Then in the following two assertions, for  $0 < \alpha \leq 1$ , (a) implies (b).*

(a)

$$\|(1 + S_t)^{-1} - (1 + C)^{-1}\| = O(t^\alpha), \quad t \downarrow 0. \quad (2.1)$$

(b) *For any  $\delta > 0$  with  $0 < \delta \leq 1$ ,*

$$\|F(t/n)^n - e^{-tC}\| = \delta^{-2} t^{-1+\alpha} e^{\delta t} O(n^{-\alpha}), \quad n \rightarrow \infty, \quad (2.2)$$

for all  $t > 0$ .

Therefore, for  $0 < \alpha < 1$  (resp.  $\alpha = 1$ ), the convergence in (2.2) is uniform on each compact  $t$ -interval in the open half line  $(0, \infty)$  (resp. in the closed half line  $[0, \infty)$ ).

Moreover, if  $C$  is strictly positive, i.e.  $C \geq \eta$  for some constant  $\eta > 0$ , the error bound on the right-hand side of (2.2) can also be replaced by  $(1 + 2/\eta)^2 t^{-1+\alpha} O(n^{-\alpha})$ , so that, for  $0 < \alpha < 1$  (resp.  $\alpha = 1$ ), the convergence in (2.2) is uniform on the closed half line  $[T, \infty)$  for every fixed  $T > 0$  (resp. on the whole closed half line  $[0, \infty)$ ).

*Sketch of Proof of Lemma.*

Put

$$F(t/n)^n - e^{-tC} = (F(t/n)^n - e^{-tS_{t/n}}) + (e^{-tS_{t/n}} - e^{-tC}).$$

For the first term on the right we have by the spectral theorem

$$\|F(t/n)^n - e^{-tS_{t/n}}\| = \|F(t/n)^n - e^{-n(1-F(t/n))}\| \leq e^{-1} n^{-1},$$

$$0 \leq e^{-n(1-\lambda)} - \lambda^n \leq e^{-1}/n, \quad \text{for } 0 \leq \lambda \leq 1.$$

For the second term, we use

$$\begin{aligned} & (1 + S_\varepsilon)^{-1} [e^{-t(\delta+S_\varepsilon)} - e^{-t(\delta+C)}] (1 + C)^{-1} \\ &= \int_0^t e^{-(t-s)(\delta+S_\varepsilon)} [(1 + S_\varepsilon)^{-1} - (1 + C)^{-1}] e^{-s(\delta+C)} ds \\ &= \int_0^{t/2} + \int_{t/2}^t. \end{aligned}$$

where  $0 < \delta \leq 1$  and  $\varepsilon > 0$ , to bound these two integrals on the right by  $(\delta^2 t)^{-1} e^{\delta t} O(\varepsilon^\alpha)$ . Taking  $\varepsilon = t/n$ , we have

$$\|e^{-tS_{t/n}} - e^{-tC}\| \leq (\delta^2 t)^{-1} e^{\delta t} O((t/n)^\alpha) = \delta^{-2} t^{-1+\alpha} e^{\delta t} O(n^{-\alpha}).$$

*Sketch of Proof of Theorems 1.1 and 1.2.*

First note that since  $C = A + B$  is itself selfadjoint and so a closed operator, by the closed graph theorem there exists a constant  $a$  such that

$$\|(1 + A)u\| + \|(1 + B)u\| \leq a\|(1 + C)u\|, \quad u \in D[C] = D[A] \cap D[B].$$

The proof of the theorem is divided into two cases, (a) the symmetric product case

$$F(t) = e^{-tB/2} e^{-tA} e^{-tB/2}, \quad (2.3)$$

and (b) the non-symmetric product case

$$G(t) = e^{-tA} e^{-tB}. \quad (2.4)$$

(a) In the symmetric case we put

$$S_t = t^{-1}(1 - F(t)) = t^{-1}(1 - e^{-tB/2} e^{-tA} e^{-tB/2})$$

and use Lemma to show that

$$\|(1 + S_t)^{-1} - (1 + C)^{-1}\| = O(t^{1/2}), \quad t \downarrow 0.$$

Put

$$A_t = t^{-1}(1 - e^{-tA}), \quad B_t = t^{-1}(1 - e^{-tB}), \quad C_t = t^{-1}(1 - e^{-tC}).$$

We have

$$\begin{aligned} 1 + S_t &= 1 + A_t + B_{t/2} - \frac{t}{4} B_{t/2}^2 + \frac{t^2}{4} B_{t/2} A_t B_{t/2} - \frac{t}{2} (A_t B_{t/2} + B_{t/2} A_t) \\ &= K_t^{1/2} (1 + Q_t) K_t^{1/2}, \end{aligned}$$

$$K_t = 1 + A_t + B_{t/2} - \frac{t}{4}B_{t/2}^2 \geq 1,$$

$$Q_t = \frac{t^2}{4}K_t^{-1/2}B_{t/2}A_tB_{t/2}K_t^{-1/2} - \frac{t}{2}K_t^{-1/2}(A_tB_{t/2} + B_{t/2}A_t)K_t^{-1/2}.$$

Then we can show

$$\|(1 + Q_t)^{-1}\| \leq 2/(3 - \sqrt{5}), \quad (2.5)$$

$$\|(1 + S_t)^{-1}K_t^{1/2}\| = \|K_t^{-1/2}(1 + Q_t)^{-1}\| \leq 2/(3 - \sqrt{5}). \quad (2.6)$$

Then we have

$$\begin{aligned} & (1 + S_t)^{-1} - (1 + C)^{-1} \\ &= (1 + S_t)^{-1}[A + B - (A_t + B_{t/2} - \frac{t}{4}B_{t/2}(1 - tA_t)B_{t/2} \\ & \quad - \frac{t}{2}(A_tB_{t/2} + B_{t/2}A_t))](1 + C)^{-1} \\ &= (1 + S_t)^{-1}(A - A_t)(1 + C)^{-1} + (1 + S_t)^{-1}(B - B_{t/2})(1 + C)^{-1} \\ & \quad + (1 + S_t)^{-1}[\frac{t}{4}B_{t/2}(1 - tA_t)B_{t/2} + \frac{t}{2}(A_tB_{t/2} + B_{t/2}A_t)](1 + C)^{-1} \\ &\equiv R_1(t) + R_2(t) + R_3(t). \end{aligned} \quad (2.7)$$

We can show the bounds

$$\|R_i(t)\| \leq ct^{1/2}, \quad i = 1, 2, 3, \quad (2.8)$$

with some constant  $c > 0$ . For instance, we can get the bound for  $R_1(t)$ , via the expression

$$\begin{aligned} R_1(t) &= [(1 + S_t)^{-1}K_t^{1/2}][K_t^{-1/2}(1 + A_t)^{1/2}] \\ &\quad \times [(1 + A_t)^{-1/2} - (1 + A_t)^{1/2}(1 + A)^{-1}](1 + A)(1 + C)^{-1} \end{aligned}$$

by (2.6) and the spectral theorem

$$\|R_1(t)\| \leq \frac{2}{3-\sqrt{5}}a\|(1 + A_t)^{-1/2} - (1 + A_t)^{1/2}(1 + A)^{-1}\| \leq ct^{1/2}.$$

(b) The non-symmetric case will follow from the symmetric case. We use the commutator argument to observe that

$$\begin{aligned} \|G(t/n)^n - F(t/n)^n\| &= \|(e^{-tA/n}e^{-tB/n})^n - (e^{-tB/2n}e^{-tA/n}e^{-tB/2n})^n\| \\ &= O(1/n). \end{aligned}$$

### 3. The Final Result

In a recent preprint [14], we have shown that if  $\kappa = 2$ , then Theorem 1.2 holds with optimal error bound  $O(n^{-1})$ . Further, the convergence is uniform on each compact  $t$ -interval in the closed half line  $[0, \infty)$ , and further, if  $C$  is strictly positive, uniform on the whole closed half line  $[0, \infty)$ .

The idea of proof is simply to iterate the resolvent equation of the first identity in (2.5) with help of its adjoint form to get

$$\begin{aligned} & (1 + S_t)^{-1} - (1 + C)^{-1} \\ &= ((1 + C)^{-1} + [(1 + S_t)^{-1} - (1 + C)^{-1}](C - S_t)(1 + C)^{-1} \\ &= (1 + C)^{-1}(C - S_t)(1 + C)^{-1} + [(C - S_t)(1 + C)^{-1}]^*(1 + S_t)^{-1}(C - S_t)(1 + C)^{-1} \\ &\equiv R'_1(t) + R'_2(t). \end{aligned}$$

Then by the same arguments together with (2.6) we can show the bounds

$$\|R'_i(t)\| = O(t), \quad i = 1, 2.$$

Therefore it turns out that the product formula (1.2) in Theorem 1.1 holds, now with ultimate error bound  $O(n^{-1})$ , properly extending and containing all the known previous related results.

Finally, we comment about optimality of the error bound  $O(n^{-1})$ . We know that if both  $A$  and  $B$  are bounded operators, then we have, in the symmetric product case (2.3),  $\|F(t/n)^n - e^{-tC}\| = O(n^{-2})$ , while, in the non-symmetric product case (2.4),  $\|G(t/n)^n - e^{-tC}\| = O(n^{-1})$ . But also in the symmetric product case, we can give an example of two unbounded selfadjoint operators  $A$  and  $B$  whose operator sum  $C = A+B$  is selfadjoint on  $D[A] \cap D[B]$  such that  $\|F(t/n)^n - e^{-tC}\| \geq L(t)n^{-1}$ , with a positive continuous function  $L(t)$  of  $t > 0$  independent of  $n$ .

Part of the present results also was briefly announced in [13].

### References

- [1] S. Deombes and B. O. Dia: *An operator-theoretic proof of an estimate on the transfer operator*, J. Functional Analysis, **165**, 240–257 (1999)
- [2] B. O. Dia and M. Schatzman: *An estimate of the Kac transfer operator*, J. Functional Analysis, **145**, 108–135 (1997)
- [3] A. Doumeki, T. Ichinose and Hideo Tamura: *Error bound on exponential product formulas for Schrödinger operators*, J. Math. Soc. Japan, **50**, 359–377 (1998)
- [4] D. Guibourg: *Inégalités maximales pour l'opérateur de Schrödinger*, C. R. Acad. Sci. Paris, **316**, Série I Math., 249–252 (1993)
- [5] B. Helffer: *Around the transfer operator and the Trotter–Kato formula*, Operator Theory: Advances and Appl., **78**, 161–174 (1995)
- [6] T. Ichinose and S. Takanobu: *Estimate of the difference between the Kac operator and the Schrödinger semigroup*, Commun. Math. Phys., **186**, 167–197 (1997)

- [7] T. Ichinose and S. Takanobu: *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup: A unified approach to the nonrelativistic and relativistic cases*, Nagoya Math. J., **149**, 53–81 (1998)
- [8] T. Ichinose and S. Takanobu: *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup II: The general case including the relativistic case*, Electronic Journal of Probability, **5**, Paper no. 5, 1–47 (2000)
- [9] T. Ichinose and Hideo Tamura: *Error estimates in operator norm for Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **27**, 195–207 (1997)
- [10] T. Ichinose and Hideo Tamura, *Error bound in trace norm for Trotter–Kato product formula of Gibbs semigroups*, Asymptotic Analysis, **17**, 239–266 (1998)
- [11] T. Ichinose and Hideo Tamura: *Error estimates in operator norm of exponential product formulas for propagators of parabolic evolution equations*, Osaka J. Math., **35**, 751–770 (1998)
- [12] T. Ichinose and Hideo Tamura: *The norm convergence of the Trotter–Kato product formula with error bound*, to appear in Commun. Math. Phys.
- [13] T. Ichinose and Hideo Tamura: *On the norm convergence of the Trotter–Kato product formula with error bound*, to appear in Proceedings of Partial Differential Equations, Clausthal, July 24–28, 2000
- [14] T. Ichinose, Hideo Tamura, Hiroshi Tamura and V. A. Zagrebnov: *Note on the paper “The norm convergence of the Trotter–Kato product formula with error bound” by Ichinose and Tamura*, to appear
- [15] T. Kato: *Trotter’s product formula for an arbitrary pair of self-adjoint contraction semigroups*, In: Topics in Functional Analysis (G.C. Rota, ed.), Academic Press, New York, 1978, pp.185–195
- [16] H. Neidhardt and V. Zagrebnov: *On error estimates for the Trotter–Kato product formula*, Lett. Math. Phys., **44**, 169–186 (1998)
- [17] H. Neidhardt and V. Zagrebnov: *Fractional powers of selfadjoint operators and Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **35**, 209–231 (1999)
- [18] H. Neidhardt and V. Zagrebnov: *Trotter–Kato product formula and operator-norm convergence*, Commun. Math. Phys., **205**, 129–159 (1999)
- [19] M. Reed and B. Simon: *Methods of Modern Mathematical Physics I: Functional Analysis*, Revised and enlarged ed., Academic Press, New York, London 1980
- [20] Dzh. L. Rogava: *Error bounds for Trotter-type formulas for self-adjoint operators*, Functional Analysis and Its Applications, **27**, 217–219 (1993)
- [21] Z. Shen:  *$L^p$  estimates for Schrödinger operators with certain potentials*, Ann. Inst. Fourier, Grenoble, **45**, 513–546 (1995); *Estimates in  $L^p$  for magnetic Schrödinger operators*, Indiana Univ. Math. J., **45**, 817–841 (1996)
- [22] Hiroshi Tamura: *A remark on operator-norm convergence of Trotter–Kato product formula*, Integr. Equat. Oper. Theory, **37**, 350–356 (2000)
- [23] H. F. Trotter: *On the product of semi-groups of operators*, Proc. Amer. Math. Soc., **10**, 545–551 (1959)